



**UAB**  
Universitat Autònoma  
de Barcelona



# Timing and Duration of Neighbourhood Disadvantage: A Life Course Perspective on Health Behaviours

---

Justė Lekštytė (UAB/CED/CEDAR)

Johan Junkka (CEDAR)

European Population Conference

June 4, 2026

# Introduction

---

Growing up in disadvantaged areas is linked to:

- Poorer health
- Weaker educational & behavioural outcomes (Brandén et al., 2023)

Previous literature is mostly *cross sectional*, does not focus on *whole childhood*, focuses on *health outcomes*.

Health behaviours strongly shape long-term health outcomes

*How do the timing and duration of exposure to neighbourhood disadvantage during childhood shape adult health behaviours?*

# Theoretical background

## Mechanisms of neighbourhood influence

- institutional-environmental
  - resource-poor environments -> limited access -> lower health literacy and skills in adulthood (Aizer & Currie, 2004; Clarke et al., 2014; Kirby & Kaneda, 2005)
- social-interactive
  - interaction with peers and adult role models -> providing information, setting norms (Beaudoin, 2009)
  - preventable diseases linked to behavior
    - neighborhoods with higher smoking prevalence -> lung cancer and other smoking-related diseases
    - peer influence to substance abuse and other risk taking behavior -> persisting life course behavioral patterns (Jang & Kim, 2023)

# Theoretical background

---


## **Cumulative exposure**

- Short exposure to disadvantage -> lower influence on health exposure  
(Vanhoutte et al., 2017)

## **Sensitive periods**

- early childhood model (Leventhal, 2018)
  - Period of primary socialisation - development of cognitive and educational skills
- adolescence model
  - children become more autonomous and explorative, making neighbourhoods more important
  - risky behaviours that peers may encourage


# Data



## Swedish register data

- 1990-1993 cohorts
- N = 398,301


# Variables



## Health behaviours:

- Behaviour-related hospitalisations (5.7%)
  - obesity (E66)
  - substance-related disorders (ICD-10 F10-19)
  - ambulatory care sensitive conditions

# Variables




## Neighbourhood exposure:

- DeSO administrative units

DeSO areas in Sweden




# Variables



## Neighbourhood exposure:

- DeSO administrative units
- Deprivation index
  - individuals aged 25–64 years:
    - low educational status
    - low income
    - unemployment
    - receipt of social welfare
- Top 10% → disadvantaged

Disadvantaged areas in Sweden 1990 (DeSO 2025)



# Variables

---

## **Time in-variant**

- Gender
- Parental migration status
- Health at birth
- Parental education at birth

## **Time varying**

- Parental income (disposable household income)
- Parental employment (0, 1, 2)
- Number of children in the household (1, 2, 3+)
- Parental union (stable, single parent)
- Parental health behaviours

# Method

- Marginal structural models estimated via inverse probability of treatment weighting
- Step 1: cumulative disadvantage
  - Never experienced disadvantage
  - 1 - 2 years
  - 3 - 9 years
  - 10 - 14 years
  - 15 - 17 years
- Step 2: sensitive periods
  - Never experienced disadvantage,
  - Only in early childhood,
  - Only during the early school years,
  - Only during adolescence,
  - Early childhood and early school years
  - Early childhood and adolescence
  - Early school years and adolescence
  - In three periods of childhood.

# Results






Figure 1. Descriptive statistics

# Results




**Figure 2. Cumulative disadvantage and health behaviours in adulthood.** Notes: 95% confidence interval in brackets, Blue dots represent MSM model with IPTW weights with time in varying controls.

# Results



**Figure 4. Sensitive periods and health behaviours in adulthood.** Notes: 95% confidence interval in brackets, Blue dots represent MSM model with IPTW weights with time in varying controls.

# Results



**Figure 4. Sensitive periods and health behaviours in adulthood.** Notes: 95% confidence interval in brackets, Blue dots represent MSM model with IPTW weights with time in varying controls.

## Conclusions:

- Effects depend on when and for how long disadvantage is experienced
  - Longer exposure -> higher hospitalisation risk
  - Early school years/adolescence appear more sensitive than early childhood

## Future steps:

- Assessing different measures of disadvantage
- Mechanisms - does disadvantage operate through structural access to healthcare or social neighbourhood processes

**Thank you for your attention!**

Justė Lekštytė

Universitat Autònoma de Barcelona, Centro de  
Estudios Demográficos

[jlekstyte@ced.uab.es](mailto:jlekstyte@ced.uab.es)

**LIFELONGMOVE**

Understanding spatial mobility  
from early life into adulthood

**European Research Council  
Consolidator Grant (CoG)**


Ref: 101043981  
Period: Jan 2023 – Dec 2027

LIFE  
LONG  
MOVE




# Appendix

Condition	ICD-10 coding
Chronic conditions	
Anemia	D501, D508, D509
Asthma	J45, J46
Diabetes	E101–E108, E110–E118, E130–E138, E140–E148
Congestive heart failure	I50, I110, J81
Hypertension	I10, I119
Chronic obstructive pulmonary disease	J41, J42, J43, J44, J47 J20 if secondary diagnosis J41, J42, J43, J44 or J47
Angina pectoris	I20, I240, I248, I249
Acute conditions	
Bleeding ulcer	K250, K251, K252, K254, K255, K256, K260, K261, K262, K264, K265, K266, K270, K271, K272, K274, K275, K276, K280, K281, K282, K284, K285, K286
Diarrhea	E86, K522, K528, K529
Epileptic seizure	O15, G40, G41, R56
Inflammatory diseases of female pelvic organs	N70, N73, N74
Renal tubulo-interstitial disease	N390, N10, N11, N12, N136
Ear, nose and throat infection	H66, H67, J02, J03, J06, J312



**Cumulative disadvantage and health behaviours in adulthood.** *Notes: 95% confidence interval in brackets, Blue dots represent MSM model with IPTW weights with time in varying controls. Pink dots represent logistic regression without any controls. Yellow dots represent logistic regression with time in varying controls. Green dots represent logistic regression with time-varying and time-invariant controls.*



**Sensitive periods and health behaviours in adulthood.** *Notes: 95% confidence interval in brackets, Blue dots represent MSM model with IPTW weights with time in varying controls. Pink dots represent logistic regression without any controls. Yellow dots represent logistic regression with time in varying controls. Green dots represent logistic regression with time-varying and time-invariant controls.*